

This article was downloaded by: [UIC University of Illinois at Chicago]

On: 3 September 2009

Access details: Access Details: [subscription number 907003094]

Publisher Routledge

Informa Ltd Registered in England and Wales Registered Number: 1072954 Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



## Nutrition and Cancer

Publication details, including instructions for authors and subscription information:

<http://www.informaworld.com/smpp/title-content=t775653687>

### Morinda Citrifolia (Noni) Reduces Cancer Risk in Current Smokers by Decreasing Aromatic DNA Adducts

Mian-Ying Wang<sup>a</sup>; Lin Peng<sup>a</sup>; May Nawal Lutfiyya<sup>a</sup>; Eric Henley<sup>a</sup>; Vicki Weidenbacher-Hoper<sup>a</sup>; Gary Anderson<sup>a</sup>

<sup>a</sup> College of Medicine at Rockford, University of Illinois Chicago, Rockford, Illinois, USA

Online Publication Date: 01 September 2009

**To cite this Article** Wang, Mian-Ying, Peng, Lin, Lutfiyya, May Nawal, Henley, Eric, Weidenbacher-Hoper, Vicki and Anderson, Gary (2009) 'Morinda Citrifolia (Noni) Reduces Cancer Risk in Current Smokers by Decreasing Aromatic DNA Adducts', *Nutrition and Cancer*, 61:5, 634 — 639

**To link to this Article:** DOI: 10.1080/01635580902825605

**URL:** <http://dx.doi.org/10.1080/01635580902825605>

## PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article may be used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

## ***Morinda Citrifolia* (Noni) Reduces Cancer Risk in Current Smokers by Decreasing Aromatic DNA Adducts**

**Mian-Ying Wang, Lin Peng, May Nawal Lutfiyya, Eric Henley, Vicki Weidenbacher-Hoper, and Gary Anderson**

*University of Illinois Chicago, College of Medicine at Rockford, Rockford, Illinois, USA*

---

**Quantitative determination of aromatic DNA adducts in peripheral blood lymphocytes (PBLs) of current smokers is an useful surrogate biomarker for the evaluation of environmental carcinogen exposure or chemopreventive intervention. In this study, we examined the impact of Tahitian Noni Juice (TNJ) on the aromatic DNA adducts of PBLs, before and after a 1-mo intervention, using <sup>32</sup>P postlabeling assay. Of 283 enrolled, 203 smokers completed the trial. Aromatic DNA adducts levels in all participants were significantly reduced by 44.9% ( $P < 0.001$ ) after drinking 1 to 4 oz of TNJ for 1 mo. Dose-dependent analyses of aromatic DNA adduct levels showed reductions of 49.7% ( $P < 0.001$ ) in the 1-oz TNJ group and 37.6% ( $P < 0.001$ ) in the 4-oz TNJ group. Gender-specific analyses resulted in no significant differences in the 4-oz TNJ groups. Interestingly, the 1-oz TNJ group showed a reduction of 43.1% ( $P < 0.001$ ) in females compared with 56.1% ( $P < 0.001$ ) in males. The results suggest that drinking 1 to 4 oz of TNJ daily may reduce the cancer risk in heavy cigarette smokers by blocking carcinogen-DNA binding or excising DNA adducts from genomic DNA.**

---

### **INTRODUCTION**

Cigarette smoke is the single known, leading preventable cause of lung cancer. In 2006, an estimated 170,000 cancer deaths in the United States were caused by tobacco smoke. Approximately 45 million Americans are current smokers, of which nearly 11 million, or 24%, are between the ages of 18 and 44. If this current trend continues, the 2030 global death toll due to smoking will exceed 9 million (1). On average, adults who smoke cigarettes die 14 years earlier than nonsmokers.

It is known that chemical carcinogenesis is a multistage process involving initiation, promotion, and progression (2). Initiation, the critical first step in carcinogenesis, requires the binding of active chemical carcinogens to genomic DNA and forming DNA adducts, which result in mutation and consequent cancer

(3). Therefore, the detection and quantification of DNA adducts is an important, fundamental, early indicator of cancer risk (4).

Tobacco smoke contains approximately 4,700 compounds and more than 68 carcinogens. The most important carcinogens in tobacco smoke are polycyclic aromatic hydrocarbons (PAHs) such as benzo(a)pyrene (5,6). Aromatic DNA adducts have been detected in lung, breast, and pancreatic cancer tissues and their surrounding normal tissues as well as in peripheral blood lymphocytes (PBLs) (7–9). Molecular epidemiological studies have demonstrated that aromatic DNA-adduct levels in smokers are significantly higher than in nonsmokers (10,11). The smoking-related aromatic DNA adduct levels decrease on smoking cessation and are positively correlated with pack years (packages of cigarettes smoked per day  $\times$  years smoked) (12–14). Furthermore, epidemiological studies have also demonstrated that the aromatic DNA-adduct levels in PBLs are a good surrogate biomarker for lung-tissue adducts and for procarcinogenic damage (15). A recent study indicated that peripheral blood aromatic DNA-adduct levels predicted lung-cancer risk in current smokers. Male smokers with higher aromatic DNA adducts, detected by <sup>32</sup>P postlabeling assay, had a threefold increased risk of lung cancer compared with males with low aromatic adduct levels (16). Additionally, a smoking-specific <sup>32</sup>P-postlabeling DNA adduct pattern called Diagonal Radioactive Zone (DRZ) has been successfully detected in the PBLs of current smokers, which represents the complex mixture of carcinogens from cigarette smoke and the major components of the aromatic DNA adducts in smokers (17). Meta-analysis has revealed a significant association between the elevated, tobacco smoke-induced, aromatic DNA-adduct levels and cancer risk (18). Thus, measuring smoking-related aromatic DNA adducts may serve as a tool for monitoring cigarette smoke exposure, assessing subsequent risk and the efficacy of chemopreventive strategies (19–24). The variance in aromatic DNA adduct patterns and levels in the PBLs of current smokers before and after preventive intervention can act as a surrogate biomarker for evaluating the ability of preventive agents to reduce PAH carcinogen-induced DNA damage, reducing cancer risk.

*Morinda citrifolia* (Noni) has been used by Polynesians for over 2,000 years as a tropical folk medicine. It is reported to

---

Submitted 20 July 2008; accepted in final form 2 February, 2009.  
Address correspondence to Mian-Ying Wang, Department of Pathology, University of Illinois-Chicago, College of Medicine at Rockford, Rockford, IL 61107. Phone: 815-395-5674. Fax: 815-395-5793. E-mail: mianwang@uic.edu

have a broad range of health benefits including preventive properties. Due to its medicinal and nutritional value, it is considered the "Queen" of the other 80 species belonging to the Rubiaceae family (25). Recent scientific studies have supported the Polynesians' claim of its unusual healing power. These studies have implicated noni as a natural remedy that lowers blood pressure, reduces joint swelling, stops internal and external infection, clears up congestion, and prevents precancer cells from growing (26). It has been reported that damnacanthol from noni roots induces normal phenotypes in *ras*-transformed tumor cells (27). Also, noni fruit extract inhibits the proliferation of breast cancer cells (28). Our previous studies have demonstrated that juice made from Tahitian noni fruits (Tahitian Noni<sup>®</sup> juice or TNJ) is a strong antioxidant, anti-inflammatory, antiangiogenic, and antiproliferative nutritional supplement (29). More specifically, TNJ has been demonstrated to significantly reduce an aromatic carcinogen, dimethylbenzo(a)anthracene (DMBA), and to induce DNA adduct formation in different organs of male C57/BL6 mice and female SD rats (30). The latest study in our laboratory indicated that TNJ was able to prevent DMBA-induced mammary carcinogenesis at the initiation and postinitiation stages in female SD rats. The reduction of the DMBA adduct level in the mammary gland tissues might be 1 of the preventive mechanisms of TNJ (31).

In this study, we examined whether TNJ was able to reduce the cancer risk by decreasing the aromatic DNA adduct levels in PBLs of current smokers after 1 mo TNJ intervention. The TNJ used in this trial was donated by Morinda Holding Inc. TNJ is a blend of Noni juice, formulated with blueberry and grape juice. Noni juice is the dominate component of TNJ (32). We hypothesized that TNJ was a good chemopreventative nutritional supplement and predicted that the aromatic DNA level in PBLs of current smokers would be reduced after 1 mo of chemopreventive intervention with TNJ. Our results are very promising.

## METHODS

The trial received human subject's approval from the Institutional Review Board of University of Illinois College of Medicine at Rockford.

### Enrollment

Intervention participants for this study met the following criteria: 1) healthy adult male or female, 2) 18 to 65 yr in age, 3) smoke more than 20 cigarettes per day, 4) a smoking history exceeding 1 yr, 5) no usage of prescribed medicine or antioxidant vitamins in the last 3 mo, and 6) willingness to complete a 1-mo trial. In addition, a control group of 20 male and 22 female nonsmokers were also recruited to this study to compare the baseline level of the aromatic DNA adducts between smokers and nonsmokers.

Participants were randomly assigned to either a 1- or 4-oz TNJ group on a 1-to-1 male to female ratio. Study participants

in the 1-oz group were asked to drink their assigned TNJ dose in the morning on an empty stomach. Those in the 4-oz group were asked to drink a 2-oz dose twice daily, 1 in the morning on an empty stomach and 1 before bedtime. Additionally, all participants were asked to drink 1 cup of pure water after consuming the TNJ.

Immediately after enrollment, 10 ml of whole blood were drawn from each participant. These pretest blood samples were used to determine baseline level of the aromatic DNA-adduct levels in both current smokers and nonsmokers. From these blood samples, we were able to assess DNA damage caused by cigarette smoking compared with nonsmokers. At the conclusion of the trial, 10 ml of blood sample was again drawn from each participant. These posttest blood samples were analyzed for aromatic DNA adduct levels after the 1-mo intervention. The comparison of prearomatic and postaromatic DNA adduct level was performed in current smokers. We predicted that the reduced aromatic DNA adduct level in current smokers would be observed if TNJ was an active chemopreventive nutritional supplement.

### Preparation of PBLs and DNA Isolation

The blood samples (pretrial and posttrial) were drawn into vacationers (green top tubes from BD, CITY, STATE) containing heparin. The tubes were centrifuged at 1,500 *g* for 20 min to remove plasma. The rest part of blood was transferred into a 50 ml tube mixed with 15 ml of red blood cell lysis buffer [150 mM NH<sub>4</sub>Cl, 10 mM sodium hydrogen carbonate, 1 mM ethylenediamine tetraacetic acid (EDTA), pH 7]. The tube was then inverted several times followed by incubation at room temperature for 5 min. Samples were then centrifuged at 300 *g* for 10 min at 4°C to isolate PBLs, and the supernatant was discarded. The PBL pellet was washed with 5 ml of red blood cell lysis buffer, incubated at room temperature, and centrifuged at 300 *g* for 10 min at 4°C, and the supernatant was discarded. This procedure was repeated 3 times. The final PBLs pellet was resuspended with 0.5 ml of buffer 1 (150 mM sodium chloride and 10 mM EDTA, pH 8.0), and cells were lysed by Vortex for 2 min at full speed. DNA was then isolated by following the standard procedure of instruction (Fast DNA-kit from Q-Biogene, CITY, STATE) (33). The purified DNA samples were stored at -80°C until DNA adduct analysis was performed using <sup>32</sup>P-postlabeling assay.

### <sup>32</sup>P-Postlabeling Assay

Aromatic DNA adduct levels were measured by the nuclease P1 procedure of <sup>32</sup>P-postlabeling assay. DNA (5–10 μg) was digested with micrococcal endonuclease (Sigma Chemicals Inc, CITY, STATE) and spleen phosphodiesterase (Sigma Chemicals Inc) to 3' mononucleotides. The unmodified normal nucleotides were dephosphorylated with nuclease P1 (USBiological, CITY, STATE), and the adducted nucleotides were labeled with [ $\gamma$ -<sup>32</sup>P]adenosine-5'-triphosphate (ICN Pharmaceuticals,

Inc., Irvine, CA) and T4 polynucleotide kinase (Sigma Chemicals Inc) (34). The labeled products were purified and separated by thin layer chromatography using homemade polyethylenimine (PEI) cellulose thin layer chromatography (TLC) sheets. After the first development overnight, the chromatogram was cut into  $1.0 \times 2.4$  cm strips above the origin point and transferred to a fresh PEI-cellulose TLC sheet by a magnet transfer technique. Two-dimensional chromatography was completed in different solvents as described in previous publications (35). Adducts were detected by autoradiography and quantified by scintillation counting. Adduct levels were expressed as a relative adduct labeling value, which is a count ratio of adducted nucleotides to total nucleotides used as a standard in this assay (36).

### Preparation of the PEI Cellulose TLC Sheets

The PEI cellulose TLC plate was homemade using IONAC Corcat P-600 PEI (Sybron Chemicals Inc., Wellford, SC) and Cellulosepulver MN 301 (Cat #816250 from Macherey Nagel, Germany; Carrier Alltech Associations, Deerfield, IL). 54.0 g of cellulose plus 5% PEI 42 ml (pH 6.0) and ddH<sub>2</sub>O 270 ml to make 4 sheets by following the original procedure developed by Randerath (37). The procedures of the preparation of PEI cellulose TLC sheets and/or <sup>32</sup>P-postlabeling assay is discussed elsewhere (38).

To avoid systemic errors, the aromatic DNA adduct analysis for all of the DNA samples (before and after intervention) and DNA samples from nonsmokers were analyzed at the same time using the same batch number of reagents and the same preparation of PEI cellulose TLC sheets. The <sup>32</sup>P postlabeling assay was carried out by an expert. To overcome possible bias, all the DNA samples were coded and blinded.

### Statistical Analysis and Data Interpretation

A power analysis was performed to estimate the number of cases needed to detect significant effects (39). All coded adduct data were entered into a predesigned database by a research assistant in a blinded manner. All data analyses were completed by an experienced and independent epidemiologist. Because the study was designed to compare the predate and postdata, all analyses were conducted on the paired cases of each group. The code number and the results were not released until all data analyses were completed. To assess the modification of TNJ on the aromatic DNA adduct levels, the averaged mean of the DNA adduct levels was compared before and after the trial in each group using a paired Student's *t*-test (40).

### RESULTS

A typical aromatic DNA adduct profile from the PBLs were detected before and after 1 mo TNJ intervention (see Fig. 1). In Fig. 1, panel A represents a typical aromatic DNA adduct profile, whereas panel B shows significant TNJ-induced reductions in size and density of the aromatic DNA adduct spots 1 to 4. Additionally, the characteristic DRZ, which contained a complex mixture of aromatic and/or hydrophobic adducts with




FIG. 1. Typical profile of the blood aromatic DNA adduct pattern in a current smoker before (panel A) and after (panel B) consumption of 1-oz Tahitian Noni Juice. Numbers 1 to 4 represent adduct spots in the blood aromatic adduct patterns. The smoking-specific Diagonal Radioactive Zone (DRZs) are represented by the dotted-line areas.

a variety of chemicals, was greatly diminished in size and pigmentation. The aromatic DNA adduct level in current smokers was 3.5 times higher than nonsmokers ( $155.1 \times 10^{-9}$  in smokers vs.  $34.7 \times 10^{-9}$  in nonsmokers); this finding was consistent with other reports (41).

To further test our hypothesis that TNJ reduces aromatic DNA adducts, we analyzed the predate and postdata from the pooled 1- and 4-oz TNJ-treated groups (Table 1). This analysis showed a aromatic DNA adduct-level decrease of 44.9% ( $P < 0.001$ ) after drinking 1- to 4-oz TNJ doses daily for 1 mo. Additional stratified analyses of the combined 1- and 4-oz TNJ data suggested that the aromatic DNA adduct level in participants' PBLs were significantly reduced ( $P < 0.001$ ) by 49% in male smokers and 40.8% in female smokers, suggesting that males respond better to TNJ than females, although the difference of the reductions between male and female was not statistically significant.

Table 2 displays separate TNJ dose and gender effects on reducing aromatic DNA adduct levels among participant smokers. The 1-oz TNJ treatment produced a reduction of 49.7% among combined genders. However, when genders were evaluated separately, males showed a 56.1% reduction compared with 43.1% for the females. All of the 1-oz TNJ-induced reductions

TABLE 1  
Aromatic DNA adduct levels in current smokers before and after consuming 1 or 4 oz of TNJ<sup>a</sup>

Gender	Time	RAL $\times 10^{-9}$	Cases (n)	Changes (%)	P Value
	Before	155.1 $\pm$ 9.0	203		
	After	85.5 $\pm$ 4.7	203	44.9 $\downarrow$	<0.001
Males	Before	156.9 $\pm$ 13.4	101		
	After	80.1 $\pm$ 6.6	101	49.0 $\downarrow$	<0.001
Females	Before	153.3 $\pm$ 12.0	102		
	After	90.8 $\pm$ 6.8	102	40.8 $\downarrow$	<0.001

<sup>a</sup>Abbreviations are as follows: TNJ, Tahitian Noni Juice; RAL, relative adduct labeling.

TABLE 2  
Aromatic DNA adduct levels in different groups before and after consuming TNJ<sup>a</sup>

Gender	Trial Time	Adduct Levels ( $\times 10^{-9}$ )	
		1 oz TNJ	4 oz TNJ
	Before	161.4 $\pm$ 11.8	146.3 $\pm$ 9.0
	After	81.3 $\pm$ 6.3	91.3 $\pm$ 65.9
	Changes	49.7% reduction	37.6% reduction
	<i>N</i>	118	85
	<i>P</i> Value	<0.001	<0.001
Male	Before	162.9 $\pm$ 17.1	148.5 $\pm$ 21.8
	After	71.5 $\pm$ 8.0	92.1 $\pm$ 10.9
	Changes	56.1% reduction	38% reduction
	<i>N</i>	59	42
	<i>P</i> Value	<0.001	0.003
Female	Before	159.9 $\pm$ 16.3	144.2 $\pm$ 17.4
	After	91.3 $\pm$ 9.6	90.5 $\pm$ 9.5
	Changes	43.1% reduction	37.3% reduction
	<i>N</i>	59	43
	<i>P</i> Value	<0.001	0.002

<sup>a</sup>Abbreviation is as follows: TNJ, Tahitian Noni Juice.

were significant at  $P < 0.001$ . The 4-oz treatment produced a significant ( $P < 0.001$ ) reduction of 37.6% among combined genders; but when genders were analyzed separately, the reductions were 38% ( $P = 0.003$ ) and 37.3% ( $P = 0.002$ ) for males and females, respectively.

## DISCUSSION

This study was designed to examine whether consuming TNJ, during a 1-mo intervention, would reduce blood aromatic DNA damage in current smokers. Our analyses revealed 5 important findings. First, current heavy smokers have 3.5-fold higher aromatic DNA adduct level than nonsmokers, indicating that aromatic carcinogens in the cigarette smoke are responsible for the elevated level of DNA damage. Second, when examining pretest and posttest aromatic DNA adduct levels of pooled genders, but separate TNJ-dose levels, the data revealed that TNJ produced highly significant reductions ( $P < 0.001$ ). Third, separate data analyses by gender and TNJ dose also produced significant reductions ( $P \leq 0.003$ ) of aromatic adduct levels. Fourth, irrespective of separate or pooled genders, the 1-oz TNJ dose produced a highly significant ( $P < 0.001$ ) reduction in the aromatic DNA adduct levels. Fifth, although reductions from the 4-oz TNJ dose were also significant ( $P \leq 0.003$ ), they were less so than for the 1-oz TNJ participants ( $P < 0.001$ ).

In our study, TNJ showed a more positive impact on reducing the aromatic DNA adduct levels in male smokers than female smokers. Whether this finding is correlated with women being more prone to cigarette-related lung cancer remains unknown (42).

Individual susceptibility to DNA damage may be indicated by the higher DNA adduct levels in PBLs of current smokers. Reducing the aromatic DNA adduct levels by consumption of 1 oz of TNJ suggests that at this dose level, the efficacy may reach a plateau such that greater reduction in DNA adduct levels with higher doses of TNJ may not occur or may do so with extended use.

Additionally, the results of this study suggest that the efficacy of TNJ on the reduction of aromatic DNA adduct levels may be extremely high in current smokers. If smokers consume 1 to 4 oz of TNJ daily, their bodies might detoxify, thus reducing the harmful impact on aromatic DNA adduct levels in cigarette smoke and possibly reducing the risk of cancer.

These findings are important because the DNA adduct levels are a cancer-risk marker and as such represent the genotoxicity, or DNA damage, smoking causes (43). If TNJ consumption reduces aromatic DNA adduct levels, cancer risk may also be reduced. In addition to cancer risk reduction, DNA adduct levels are also good biomarkers for all degenerative diseases associated with smoking (44). As noted earlier, molecular epidemiologic studies have demonstrated that aromatic DNA adduct levels in smokers are significantly higher than in nonsmokers; our data has conformed this observation. Furthermore, these types of smoking-related DNA adducts decreased with cessation of smoking and were positively correlated with pack years (packages of cigarettes smoked per day  $\times$  years smoked) (45). Additionally, findings from 1 study indicated that blood-aromatic DNA adduct levels predicted lung-cancer risk in current smokers and that male smokers with higher aromatic DNA adduct levels detected by <sup>32</sup>P postlabeling assay had a threefold increased risk of lung cancer compared with males with low aromatic adduct levels (16). The mechanisms for the reduction of aromatic DNA adduct in current smokers by consuming TNJ needs further study to determine if TNJ inhibits Phase 1 enzyme activities and enhances Phase 2 and DNA repair enzyme activities (46). The impact of TNJ on gene susceptibility of current smokers will be investigated soon (47).

Finally, based on the response of study participants to a questionnaire, no adverse events were observed during the trial. Approximately 14% of the study participants reported that the number of cigarettes smoked daily was slightly reduced and they are willing to continue drinking TNJ. The study results suggest that TNJ may have a cancer preventive effect in current smokers. The significant reduction of aromatic DNA adduct level by consuming TNJ might be associated with the reduced number of cigarettes they smoked during this trial. Long-term impacts of TNJ on the aromatic DNA adduct level in current smokers needs further study.

## ACKNOWLEDGMENT

This research project was supported by a grant from Morinda Holdings, Inc., which provided all of the noni juice used in the intervention.

## REFERENCES

1. U.S. Department of Health and Human Services: *Women and Smoking—A Report of the Surgeon General*. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health, 2001.
2. Oliveira PA, Colaço A, Chaves R, Guedes-Pinto H, De-La-Cruz P LF, et al.: Chemical carcinogenesis. *An Acad Bras Cienc* **79**, 593–616.
3. Randerath E, Miller RH, Dinesh M, Avittis TA, Dunford A, et al.: Covalent DNA damage in tissues of cigarette smokers as determined by <sup>32</sup>P-postlabeling assay. *J Natl Cancer Inst* **81**, 341–347, 1989.
4. Nesnow S, Ross J, Nelson G, Holden K, Erexson G, et al.: Quantitative and temporal relationships between DNA adduct formation in target and surrogate tissues: implications for biomonitoring. *Environ Health Perspect* **101**(3 Suppl), 37–42, 1993.
5. Sepetdjian E, Shihadeh A, and Saliba NA: Measurement of 16 polycyclic aromatic hydrocarbons in narghile waterpipe tobacco smoke. *Food Chem Toxicol* **46**, 1582–1590, 2008.
6. Kriek E, Rojas M, Alexandrov K, and Bartsch H: Polycyclic aromatic hydrocarbon-DNA adducts in humans: relevance as biomarkers for exposure and cancer risk. *Mut Res* **1–2**, 215–231, 1998.
7. Li D, Wang M, Dhingra K, and Hittelman WN: Aromatic DNA adducts in adjacent tissues of breast cancer patients: clues to breast cancer etiology. *Cancer Res* **56**, 287–293, 1996.
8. Wang M, Abbruzzese JL, Friess H, Hittelman WN, Evans DB, et al.: DNA adducts in human pancreatic tissues and their potential role in carcinogenesis. *Cancer Res* **58**, 38–41, 1998.
9. Mooney LA, Santella RM, Covey L, Jeffrey AM, Bigbee W, et al.: Decline of DNA damage and other biomarkers in peripheral blood following smoking cessation. *Cancer Epidemiol Biomarkers Prev* **4**, 627–634, 1995.
10. Zhang LF, Xu K, and Li XM: Quantitative study on DNA-adduct in normal lung tissues of smoking and non-smoking lung cancer patients. *Zhonghua Yu Fang Yi Xue Za Zhi* **28**, 206–209, 1994.
11. Wiencke JK, Thurston SW, Kelsey KT, et al.: Early age at smoking initiation and tobacco carcinogen DNA damage in the lung. *J Natl Cancer Inst* **91**, 614–619, 1999.
12. Schoket B, Kostic S, and Vincze I: Determination of smoking-related DNA adducts in lung-cancer and non-cancer patients. *IARC Sci Publ* **124**, 315–319, 1993.
13. Kriek E, Van Schooten FJ, Hillebrand MJ, Van Leeuwen FE, Den Engelse L, et al.: DNA adducts as a measure of lung cancer risk in humans exposed to polycyclic aromatic hydrocarbons. *Environ Health Perspect* **99**, 71–75, 1993.
14. Wiencke JK, Kelsey KT, Varvony A, et al.: Correlation of DNA adducts in blood mononuclear cells with tobacco carcinogen-induced damage in human lung. *Cancer Res* **55**, 4910–4914, 1995.
15. Phillips DH, Hewer A, Martin CN, Garner R, and King MM: Correlation of DNA adduct levels in human lung with cigarette smoking. *Nature (Lond)* **336**, 790–792, 1988.
16. van Schooten FJ, Hillebrand MJ, van Leeuwen FE, et al.: Polycyclic aromatic hydrocarbon-DNA adducts in lung tissue from lung cancer patients. *Carcinogenesis* **11**, 1677–1681, 1990.
17. Peluso M, Neri M, Margarino, Mereu C, Munnia A, et al.: Comparison of DNA adduct levels in nasal mucosa, lymphocytes and bronchial mucosa of cigarette smokers and interaction with metabolic gene polymorphisms. *Carcinogenesis* **25**, 2459–2465, 2004.
18. Veglia F, Matullo G, and Vineis P: Bulky DNA adducts and risk of cancer: a meta-analysis. *Cancer Epidemiol Biomarkers Prev* **12**, 157–160, 2003.
19. Hafizur Rahman M, Iqbal Arslan M, Chen Y, Ali S, Parvin T, et al.: Polycyclic aromatic hydrocarbon-DNA adducts among rickshaw drivers in Dhaka City, Bangladesh. *Int Arch Occup and Environ Health* **76**, 533–538, 2003.
20. Hsu TM, Zhang YJ, and Santella RM: Immunoperoxidase quantitation of 4-aminobiphenyl- and polycyclic aromatic hydrocarbon-DNA adducts in exfoliated oral and urothelial cells of smokers and nonsmokers. *Cancer Epidemiol Biomarkers Prev* **6**, 193–199, 1997.
21. Castaño-Vinyals G, Talaska G, Rothman N, Alguacil J, Garcia-Closas M, et al.: Bulky DNA adduct formation and risk of bladder cancer. *Cancer Epidemiol Biomarkers Prev* **16**, 2155–2159, 2007.
22. Gyorfy E, Anna L, Gyori Z, et al.: DNA adducts in tumor, normal peripheral lung and bronchus, and peripheral blood lymphocytes from smoking and non-smoking lung cancer patients: correlations between tissues and detection by <sup>32</sup>P-postlabelling and immunoassay. *Carcinogenesis* **25**, 1201–1209, 2004.
23. Alexandrov K, Rojas M, and Rolando C: DNA damage by benzo(a)pyrene in human cells is increased by cigarette smoke and decreased by a filter containing rosemary extract, which lowers free radicals. *Cancer Res* **66**, 11938–11945, 2006.
24. Mooney LA, Madsen AM, Tang D, Orjuela MA, Tsai WY, et al.: Antioxidant vitamin supplementation reduces benzo(a)pyrene-DNA adducts and potential cancer risk in female smokers. *Cancer Epidemiol Biomarkers Prev* **14**, 237–242, 2005.
25. Morton J: The ocean-going noni, or Indian mulberry (*Morinda citrifolia*, Rubiaceae) and some of its “colorful” relatives. *Econ Bot* **46**, 241–257, 1992.
26. Wang MY, West BJ, Jensen CJ, et al. *Morinda citrifolia* (Noni): a literature review and recent advances in noni research. *Acta Pharmacol Sin* **23**, 1127–1141, 2002.
27. Hiramatsu T: Induction of normal phenotypes in *ras*-transformed cells by damnacanthol from *Morinda citrifolia*. *Cancer Lett* **73**, 161–166, 1993.
28. Arpornsuwan T and Punjanon, T: Tumor cell-selective antiproliferative effect of the extract from *Morinda citrifolia* fruits. *Phytother Res* **20**, 515–517, 2006.
29. Division of Cancer Prevention, NCI Research in Complementary and Alternative Medicine: Developing Effective and Efficient Treatments: Noni Juice Studied for Prevention of Breast Cancer From Dr. Wang, M-Y, UIC College of Medicine at Rockford. *NCI Annual Report on Complementary and Alternative Medicine Fiscal Year 2006*.
30. Wang MY and Su C: Cancer preventive effect of *Morinda citrifolia*. *Ann NY Acad Sci* **952**, 161–168, 2002.
31. Wang MY, Peng L, Anderson G, and Nowicki D: Breast cancer preventions with *Morinda citrifolia* Fruit. *Proceedings of the National Cancer Institute; Cancer Researchers and CAM Practitioners: Fostering Collaborations; Advancing the Science*. Sponsored by the Office of Cancer Complementary and Alternative Medicine, Bethesda, Maryland, October 21–23, 2007.
32. European Commission: Commission decision of 5 June 2003 authorizing the placing on the market of “Noni juice” as a novel food ingredient under regulation: (EC) Nr. 258/97 of the European parliament and of the council. *Eur J Cancer Prev* **12**, 144, 2003.
33. MP Biomedicals: Retrieved from [http://www.mpbio.com/product\\_info.php?products\\_id=6003500](http://www.mpbio.com/product_info.php?products_id=6003500)
34. Reddy M and Randerath K: Nuclease P1-mediated enhancement of sensitivity for structurally diverse DNA adducts. *Carcinogenesis* **7**, 1543–1551, 1986.
35. Wang MY and Lu LJW: Differential effect of gestation stage on benzo(a)pyrene-induced micronucleus formation and/or covalent DNA modifications in mice. *Cancer Res* **50**, 2146–2151, 1990.
36. Lu LJW and Wang MY: Modulation of benzo(a)pyrene-induced covalent DNA modification in adult and fetal mouse tissues by gestation stage. *Carcinogenesis* **11**, 1367–1372.
37. Randerath K, Gupta RC, and Randerath E: 3H- and 32P-derivative methods for base composition and sequence analysis of RNA. *Methods Enzymol* **65**, 638–680, 1980.
38. Phillips DH and Arlt VM: The <sup>32</sup>P-postlabeling assay for DNA adducts. *Nat Protoc* **2**, 2772–2781, 2007.

39. Brookes ST, Whitley E, Peters TJ, Mulheran PA, Egger M, et al.: Subgroup analyses in randomized controlled trials: quantifying the risks of false-positives and false-negatives. *Health Technol Assess* **5**, 1–56, 2001.
40. Leandro-Merhi VA, Diez Garcia RW, Monaco DV, and Marques de Oliveira MR: Comparative analysis of nutritional status, food intake, and length of hospitalization considering two hospitals: public versus private. *Nutr Hosp* **21**, 32–37, 2006.
41. Nia AB, Maas LM, Brouwer FMC, Kleinjans JCS, and Van Schooten FJ: Comparison between smoking-related DNA adduct analysis in induced sputum and peripheral blood lymphocytes. *Carcinogenesis* **21**, 1335–1340, 2000.
42. Mucha L, Stephenson J, Morandi N, and Dirani R: Meta-analysis of disease risk associated with smoking, by gender and intensity of smoking. *Genet Med* **3**, 279–291, 2006.
43. Weitberg AB and Corvese D: Oxygen radicals potentiate the genetic toxicity of tobacco-specific nitrosamines. *Clin Genet* **43**, 88–91, 1993.
44. De Flora S, Izzotti A, Randerath K, et al. DNA adducts and chronic degenerative disease. Pathogenetic relevance and implications in preventive medicine. *Mutat Res* **366**, 197–238, 1996.
45. Breast Cancer Family Registry, Kathleen Cuninghame Consortium for Research into Familial Breast Cancer (Australasia), and Ontario Cancer Genetics Network (Canada): Smoking and risk of breast cancer in carriers of mutations in BRCA1 or BRCA2 aged less than 50 years. *Breast Cancer Res Treat* **109**, 67–75, 2008.
46. Nia AB, Kleinjans JCS, and Van Schooten FJ: Biomonitoring of tobacco smoke carcinogenicity by dosimetry of DNA adducts and genotyping and phenotyping of biotransformational enzymes: a review on polycyclic aromatic hydrocarbons. *Biomarkers* **7**, 209–229, 2002.
47. Firozi PF, Bondy ML, Sahin AA, Chang P, Lukmanji F, et al.: Aromatic DNA adducts and polymorphisms of CYP1A1, NAT2, and GSTM1 in breast cancer. *Carcinogenesis* **23**, 301–306, 2002.